

- <sup>3</sup> Beckwith, J. R., *J. Mol. Biol.*, **8**, 427 (1964).
- <sup>4</sup> Beckwith, J. R., *Biochim. Biophys. Acta*, **76**, 162 (1963).
- <sup>5</sup> Benzer, S., and S. P. Champe, these PROCEEDINGS, **48**, 1114 (1962).
- <sup>6</sup> Yanofsky, C., and P. St. Lawrence, *Ann. Rev. Microbiol.*, **14**, 311 (1960); Garen, A., and O. Siddiqi, these PROCEEDINGS, **48**, 1121 (1962); Sarabhai, A. S., A. O. W. Stretton, S. Brenner, and A. Bolle, *Nature*, **201**, 13 (1964).
- <sup>7</sup> Stent, G., *Science*, **144**, 816 (1964).
- <sup>8</sup> Franklin, N. C., and S. E. Luria, *Virology*, **15**, 299 (1961); Ames, B. N., and P. Hartman, in *Synthesis and Structure of Macromolecules*, Cold Spring Harbor Symposia on Quantitative Biology, vol. 28 (1963), p. 349.
- <sup>9</sup> Crick, F. H. C., L. Barnett, S. Brenner, and R. J. Watts-Tobin, *Nature*, **192**, 1227 (1961).
- <sup>10</sup> Pardee, A. B., F. Jacob, and J. Monod, *J. Mol. Biol.*, **1**, 165 (1959).
- <sup>11</sup> Strain 2320 and both of its derivatives had reacquired the  $B_1^+$  phenotype before they were made available to us.
- <sup>12</sup> Luria, S. E., J. N. Adams, and R. C. Ting, *Virology*, **12**, 348 (1960).
- <sup>13</sup> Weigle, J., these PROCEEDINGS, **39**, 628 (1953).
- <sup>14</sup> Orgel, A., and S. Brenner, *J. Mol. Biol.*, **3**, 762 (1961).
- <sup>15</sup> Edgar, R. S., and I. Lielausis, *Genetics*, **49**, 649 (1964).
- <sup>16</sup> Benzer, S., these PROCEEDINGS, **45**, 1607 (1959).
- <sup>17</sup> Champe, S. P., and S. Benzer, *J. Mol. Biol.*, **4**, 288 (1962).
- <sup>18</sup> Epstein, R. H., *et al.*, in *Synthesis and Structure of Macromolecules*, Cold Spring Harbor Symposia on Quantitative Biology, vol. 28 (1963), p. 375. Mutants  $rGU14$  and  $rGU74$  are suppressed by strains  $CR63(\lambda)$  and  $K12(\lambda)$ .
- <sup>19</sup> Benzer, S., these PROCEEDINGS, **47**, 403 (1961).
- <sup>20</sup> Champe, S. P., and S. Benzer, these PROCEEDINGS, **48**, 532 (1962); N24 is not a member of ambivalent classes I, II, or III.
- <sup>21</sup> Benzer, S., and S. P. Champe, these PROCEEDINGS, **47**, 1025 (1961).
- <sup>22</sup> Since the burst size of even the  $r638$  deletion mutant was increased to about the same extent as the other non-*osus* mutants, we conjecture that such "nonspecific suppression" is the result of a less stringent requirement, in 2320-R8( $\lambda$ ), for the function of the  $rII$  cistrons. The basis of this effect has not been investigated further.
- <sup>23</sup> Freese, E., these PROCEEDINGS, **45**, 622 (1959).
- <sup>24</sup> Freese, E., E. Bautz, and E. Bautz-Freese, these PROCEEDINGS, **47**, 845 (1961).
- <sup>25</sup> Whereas all 2AP-revertible nonsense mutants (Benzer and Champe<sup>5</sup>) tested are 5FU-suppressible,<sup>20</sup> none were so out of 8 2AP-revertible missense mutants tested.

---

### ERRATUM

In the article entitled "Decomposition of Vector Measures," by M. M. Rao, which appeared in the May issue of Volume 51 (1964), pages 771-774, Theorem 1 (i) should read: "(i)  $|\mu|$  is an outer measure and  $\nu$  is additive and  $s$ -bounded," etc.